

CCL17. These results suggest that M2 macrophages stimulated by RANKL/RANK signaling might recruit Tregs into the tumor microenvironment of EMPD, which thereby induces an immunosuppressive microenvironment in EMPD.

Gordon and Martinez (2010) classified the development of monocytes into mature and fully activated macrophages into three successive stages. During the third phase of activation, macrophages reach a mature functional phenotype in response to microbial and opsonic stimuli such as antibody complexes. In the present study, macrophages induced by M-CSF and IL-4 did not produce a substantial amount of CCL17; however, when these macrophages were stimulated by sRANKL, they produced CCL17 in an sRANKL dose-dependent manner. These results suggest that sRANKL may be another stimulus during the third phase of macrophage activation.

Denosumab, a fully human monoclonal antibody for RANKL, is used clinically to treat metastatic bone tumors (Azim and Azim, 2013). Our present study suggests that RANKL targeting with denosumab can be used in conjunction with the therapeutic elimination of primary EMPD to prevent local immunosuppression and metastatic disease.

CONFLICT OF INTEREST

The authors state no conflict of interest.

ACKNOWLEDGMENTS

This study was supported in part by grants-in-aid for scientific research from the Japan Society for the Promotion of Science (23791249 and 25461682).

Taku Fujimura^{1,2}, Yumi Kambayashi^{1,2}, Sadanori Furudate¹, Masayuki Asano¹, Aya Kakizaki¹ and Setsuya Aiba¹

¹Department of Dermatology, Tohoku University Graduate School of Medicine, Sendai, Japan

E-mail: tfujimura1@mac.com

²These authors contributed equally to this work.

SUPPLEMENTARY MATERIAL

Supplementary material is linked to the online version of the paper at <http://www.nature.com/jid>

REFERENCES

- Anderson DM, Maraskovsky E, Billingsley WL *et al.* (1997) A homologue of the TNF receptor and its ligand enhance T-cell growth and dendritic-cell function. *Nature* 390:175–9
- Azim H, Azim HA Jr (2013) Targeting RANKL in breast cancer: bone metastasis and beyond. *Expert Rev Anticancer Ther* 13:195–201
- Fujimura T, Furudate S, Kambayashi Y *et al.* (2013) Potential use of bisphosphonates in invasive extramammary Paget's disease: an immunohistochemical investigation. *Clin Dev Immunol* 2013:164982
- Fujimura T, Kambayashi Y, Hidaka T *et al.* (2012) Comparison of Foxp3+ regulatory T cells and CD163+ macrophages in invasive and non-invasive extramammary Paget's disease. *Acta Derm Venereol* 92:625–8
- Gordon S, Martinez FO (2010) Alternative activation of macrophages: mechanism and functions. *Immunity* 32:593–604
- Hao NB, Lu MH, Fan YH *et al.* (2012) Macrophages in tumor microenvironments and the progression of tumors. *Clin Dev Immunol* 2012:948098
- Kambayashi Y, Fujimura T, Furudate S *et al.* (2015) The possible interaction between receptor activator of nuclear factor kappa-B ligand (RANKL) expressed by extramammary Paget cells and its ligand on dermal macrophages. *J Invest Dermatol* (in press)
- Loser K, Mehling A, Loeser S *et al.* (2006) Epidermal RANKL controls regulatory T-cell numbers via activation of dendritic cells. *Nat Med* 12:1372–9
- Martinez FO, Gordon S, Locati M *et al.* (2006) Transcriptional profiling of the human monocyte-to-macrophage differentiation and polarization: new molecules and patterns of gene expression. *J Immunol* 177:7303–11
- Mantovani A, Sozzani S, Locati M *et al.* (2002) Macrophage polarization: tumor-associated macrophages as a paradigm for polarized M2 mononuclear phagocytes. *Trends Immunol* 23:549–55
- Wong BR, Josien R, Lee SY *et al.* (1997) TRANCE (tumor necrosis factor [TNF]-related activation-induced cytokine), a new TNF family member predominantly expressed in T cells, is a dendritic cell-specific survival factor. *J Exp Med* 186:2075–80
- Yasuda H, Shima N, Nakagawa N *et al.* (1998) Osteoclast differentiation factor is a ligand for osteoprotegerin/osteoclastogenesis-inhibitory factor and is identical to TRANCE/RANKL. *Proc Natl Acad Sci USA* 95:3597–602

Cathelicidin LL-37 Induces Semaphorin 3A Expression in Human Epidermal Keratinocytes: Implications for Possible Application to Pruritus

Journal of Investigative Dermatology (2015) **135**, 2887–2890; doi:10.1038/jid.2015.243; published online 16 July 2015

TO THE EDITOR

Higher density of epidermal nerve fibers has been associated with itch sensitization in the periphery, with nerve fiber density being higher in atopic dermatitis (AD) lesional skin than in normal skin (Tominaga and Takamori, 2014). Concomitantly, the epidermal levels of the

nerve repulsion factor semaphorin 3A (Sema3A) are lower in AD patients and AD model NC/Nga mice than in controls (Tominaga *et al.*, 2008). Sema3A replacement in AD model NC/Nga mice normalized hyperinnervation, resulting in the suppression of itching (Yamaguchi *et al.*, 2008; Negi *et al.*, 2012).

To defend against pathogens, human skin contains antimicrobial peptides, including cathelicidins and β -defensins, which are induced by inflammation (Schauber and Gallo, 2008; Niyonsaba *et al.*, 2009). The induction of these peptides, such as cathelicidin LL-37 and human β -defensins, produced by epidermal keratinocytes is impaired in lesional skin in patients with AD, explaining their frequent infections (Ong *et al.*, 2002; de Jongh *et al.*,

Abbreviations: AD, atopic dermatitis; NHEK, normal human epidermal keratinocyte; P2X₇R, P2X₇ receptor; Sema3A, semaphorin 3A

Accepted article preview online 29 June 2015; published online 16 July 2015




Figure 1. Effects of antimicrobial peptides on Sema3A expression in cultured NHEKs. (a) NHEKs were incubated with 10 µg ml⁻¹ hBD-1-4, LL-37, or medium alone (vehicle) for 3, 24, or 48 hours, and Sema3A mRNA expression was assayed by quantitative real-time PCR. Sema3A mRNA levels were normalized relative to those of RPS18. (b) Effects of LL-37 concentration on Sema3A mRNA expression. Values represent fold changes compared with vehicle. LL-37 induced Sema3A expression dose- and time dependently. (c) NHEKs were incubated with LL-37 for 48 hours, and the concentrations of Sema3A in the cell-free supernatants were determined by ELISA. **P* < 0.05, ****P* < 0.0001 (two-way ANOVA followed by Dunnett's test), ###*P* < 0.0001 (two-tailed Student's *t*-test) compared with vehicle. All values represent the mean ± SD of 3–6 experiments. ANOVA, analysis of variance; hBD, human β defensin; NHEK, normal human epidermal keratinocyte; RPS18, ribosome protein S18; Sema3A, semaphorin 3A.

2005). However, the interrelationships among Sema3A and antimicrobial peptides are unclear. We therefore investigated the effects of antimicrobial peptides on Sema3A expression in cultured normal human epidermal keratinocytes (NHEKs), as well as the signaling pathways involved in LL-37-induced Sema3A expression.

NHEKs derived from adult epidermis (Lonza, Basel, Switzerland) were

cultured in KBM-Gold (Lonza), a serum-free medium containing a low concentration (0.15 mM) of calcium, at 37 °C with 5% CO₂ and used within three passages. LL-37 (L¹LGDFFRKSKEKIGKEFKRIVQRIKDFLRNLPRTES³⁷) was synthesized by the solid phase method on a peptide synthesizer (model PSSM-8; Shimadzu, Kyoto, Japan). NHEKs were incubated with human β-defensin-1–4 (Peptide Institute, Osaka,

Japan) or LL-37, and the levels of Sema3A mRNA expression at different time points were assessed by quantitative real-time PCR using SYBR Premix Ex Taq (TaKaRa, Kyoto, Japan) and the primers 5'-ACCCAATATCAATGGGTG CCTTA-3' (forward) and 5'-AACACTG GATTGTACATGGCTGGA-3' (reverse). As controls, ribosome protein S18 mRNA was amplified using the primers 5'-TTTGGCGAGTACTCAACACCAACATC-3' (forward) and 5'-GAGCATATCTTCGGC CCACAC-3' (reverse).

Incubation with LL-37 for 48 hours increased Sema3A expression, whereas the individual human β-defensins had little effect (Figure 1a). This LL-37-induced Sema3A expression was dose- and time dependent (Figure 1b). Furthermore, Sema3A protein levels measured using ELISA kits (USCN Life Science, Wuhan, China) were markedly higher in the supernatants of NHEKs incubated with LL-37 than in control supernatants (Figure 1c).

LL-37 has been shown to activate rat mast cells via G protein-coupled receptors (Niyonsaba *et al.*, 2001). Therefore, the effects of cholera toxin (Wako Pure Chemical Industries, Ltd. Osaka, Japan) and pertussis toxin (Sigma-Aldrich, St Louis, MO), inhibitors of the G_s and G_i subfamilies of G protein α-subunit, respectively, on Sema3A induction in cultured NHEKs were analyzed. LL-37-induced Sema3A expression was completely inhibited by pretreatment with pertussis toxin but not cholera toxin (Figure 2a).

Activation of either mitogen-activated protein kinase or phosphatidylinositol 3 kinase was found to be involved in the LL-37 signaling pathway (Niyonsaba *et al.*, 2010). We therefore assessed the effects of mitogen-activated protein kinase and phosphatidylinositol 3 kinase inhibitors on Sema3A induction in cultured NHEKs. LL-37-induced Sema3A expression was inhibited by pretreatment with the extracellular signal-regulated kinase 1/2 inhibitor PD98059 (Cell Signaling Technology, Beverly, MA), but not the phosphatidylinositol 3 kinase inhibitor wortmannin (Sigma-Aldrich), the p38 inhibitor SB203580 (Sigma-Aldrich), and the c-Jun N-terminal kinase inhibitor SP600125 (Calbiochem, Darmstadt, Germany; Figure 2b).




Figure 2. Analysis of the signaling pathway involved in LL-37-induced Sema3A expression. NHEKs were pretreated with 100 $\mu\text{g ml}^{-1}$ cholera toxin (CT) (a), 200 ng ml^{-1} pertussis toxin (PT) (a), 20 μM wortmannin (Wort.) (b), 10 μM SB203580 (SB) (b), 10 μM PD98059 (PD) (b), 10 μM SP600125 (SP) (b), 1 μM WRW4 (c), 1 μM BBG (c), or vehicle (–) for 1 hour, followed by incubation for 24 hours with 20 $\mu\text{g ml}^{-1}$ LL-37, and Sema3A expression was assayed by quantitative real-time PCR. Values represent fold changes compared with vehicle and the mean \pm SD of 4–6 experiments. ** $P < 0.01$ and *** $P < 0.001$ compared with vehicle, ## $P < 0.01$ compared with LL-37 alone (one-way ANOVA followed by Dunnett's test). (d) Schematic diagram of the signaling pathway of LL-37-induced Sema3A expression in NHEKs. ANOVA, analysis of variance; ERK, extracellular signal-regulated kinase; JNK, c-Jun N-terminal kinase; NHEK, normal human epidermal keratinocyte; PI3K, phosphatidylinositol 3 kinase; Sema3A, semaphorin 3A.

We next examined whether formyl peptide receptor-like 1 and P2X₇ receptor (P2X₇R), both candidate LL-37 receptors (Niyonsaba *et al.*, 2009), are involved in Sema3A induction in cultured NHEKs. LL-37-induced Sema3A expression was partially inhibited by pretreatment with the P2X₇R antagonist BBG (Tokyo Chemical Industry, Tokyo, Japan) but not by pretreatment with the formyl peptide receptor-like 1 antagonist WRW4 (ABGENT, San Diego, CA; Figure 2c).

Taken together, these findings showed that the antimicrobial peptide LL-37 markedly upregulated the expression of Sema3A mRNA and protein in cultured human keratinocytes, whereas the four human β -defensins were inactive (Figure 1), indicating that LL-37 is an endogenous inducer of Sema3A

expression in keratinocytes. Vitamin D3, which increases cathelicidin expression in keratinocytes (Schauber and Gallo, 2008), might therefore also drive the LL-37-Sema3A pathway.

We found that certain G_i-coupled receptors are involved in Sema3A induction (Figure 2a). Although LL-37 has been shown to activate G protein-coupled receptors, such as formyl peptide receptor-like 1, P2X₇R, and Mas-related G protein-coupled receptor X2 (MrgX2; Barlow *et al.*, 2014), LL-37 receptors are less well-characterized in human keratinocytes. We showed that LL-37-induced Sema3A expression was partly inhibited by a P2X₇R antagonist, but not by a formyl peptide receptor-like 1 antagonist (Figure 2c), and that MrgX2 mRNA expression was undetectable in NHEKs (Umehara *et al.*,

unpublished observation). These findings suggest that, although other receptors may be involved in induction, LL-37 activation of P2X₇R is at least partly involved in Sema3A induction in keratinocytes. We also found that LL-37-induced Sema3A expression was completely inhibited by PD98059 (Figure 2b), indicating that extracellular signal-regulated kinase 1/2 signaling may be required for Sema3A induction. In contrast, SP600125 increased rather than inhibiting Sema3A expression (Figure 2b), suggesting that c-Jun N-terminal kinase signaling is involved in the suppression of Sema3A expression in keratinocytes. Altered c-Jun N-terminal kinase signaling might reduce Sema3A production in AD. Taken together, these findings suggest that LL-37 may bind to certain G_i-

coupled receptors, including P2X₇R, activating the extracellular signal-regulated kinase 1/2 signaling pathway, and finally inducing Sema3A expression in human epidermal keratinocytes (Figure 2d).

Although 70–90% of patients with psoriasis also have pruritus, they are rarely accompanied by intense itch as in AD (Yosipovitch et al., 2000; Reich and Szepletowski, 2007). We previously showed that Sema3A production was not reduced in the epidermis of psoriatic patients with or without itch (Taneda et al., 2011). In contrast to AD, high levels of expression of several antimicrobial peptides including LL-37 were observed in psoriatic lesions (Ong et al., 2002; Nomura et al., 2003), suggesting that epidermal Sema3A expression is not reduced in these patients. These findings suggest that LL-37 may restore Sema3A production in keratinocytes of certain pathological conditions such as psoriasis and that the decreased epidermal Sema3A production in AD lesional skin may be partly caused by a breakdown in this recovery system. Thus, the topical application of LL-37, which not only has antimicrobial activity but also enhances Sema3A production, might be a useful therapeutic strategy for AD.

CONFLICT OF INTEREST

The authors state no conflict of interest.

ACKNOWLEDGMENTS

This work was partly supported by KAKENHI (Grant number 26860898) and by a grant of Strategic Research Foundation Grant-aided Project for Private Universities from MEXT (Grant number S1311011).

Yoshie Umehara¹, Yayoi Kamata¹, Mitsutoshi Tominaga¹, François Niyonsaba^{2,3}, Hideoki Ogawa^{1,2} and Kenji Takamori^{1,4}

¹Institute for Environmental and Gender Specific Medicine, Juntendo University Graduate School of Medicine, Chiba, Japan; ²Atopy (Allergy) Research Center, Juntendo University Graduate School of Medicine, Tokyo, Japan; ³Faculty of International Liberal Arts, Juntendo University, Tokyo, Japan and ⁴Department of Dermatology, Juntendo University Urayasu Hospital, Urayasu, Chiba, Japan
E-mail: ktakamori@juntendo.ac.jp

REFERENCES

- Barlow PG, Findlay EG, Currie SM et al. (2014) Antiviral potential of cathelicidins. *Future Microbiol* 9:55–73
- de Jongh GJ, Zeeuwen PL, Kucharekova M et al. (2005) High expression levels of keratinocyte antimicrobial proteins in psoriasis compared with atopic dermatitis. *J Invest Dermatol* 125:1163–73
- Negi O, Tominaga M, Tengara S et al. (2012) Topically applied semaphorin 3A ointment inhibits scratching behavior and improves skin inflammation in NC/Nga mice with atopic dermatitis. *J Dermatol Sci* 66:37–43
- Niyonsaba F, Someya A, Hirata M et al. (2001) Evaluation of the effects of peptide antibiotics human β -defensins-1/2 and LL-37 on histamine release and prostaglandin D₂ production from mast cells. *Eur J Immunol* 31:1066–75

- Niyonsaba F, Nagaoka I, Ogawa H et al. (2009) Multifunctional antimicrobial proteins and peptides: natural activators of immune systems. *Curr Pharm Des* 15:2393–413
- Niyonsaba F, Ushio H, Hara M et al. (2010) Antimicrobial peptides human β -defensins and cathelicidin LL-37 induce the secretion of a pruritogenic cytokine IL-31 by human mast cells. *J Immunol* 184:3526–34
- Nomura I, Goleva E, Howell MD et al. (2003) Cytokine milieu of atopic dermatitis, as compared to psoriasis, skin prevents induction of innate immune response genes. *J Immunol* 171:3262–9
- Ong PY, Ohtake T, Blandt C et al. (2002) Endogenous antimicrobial peptides and skin infections in atopic dermatitis. *N Engl J Med* 347:1151–60
- Reich A, Szepletowski JC (2007) Mediators of pruritus in psoriasis. *Mediators Inflamm* 2007: 64727
- Schauber J, Gallo RL (2008) Antimicrobial peptides and the skin immune defense system. *J Allergy Clin Immunol* 122:261–6
- Taneda K, Tominaga M, Negi O et al. (2011) Evaluation of epidermal nerve density and opioid receptor levels in psoriatic itch. *Br J Dermatol* 165:277–84
- Tominaga M, Ogawa H, Takamori K (2008) Decreased production of semaphorin 3A in the lesional skin of atopic dermatitis. *Br J Dermatol* 158:842–4
- Tominaga M, Takamori K (2014) Itch and nerve fibers with special reference to atopic dermatitis: therapeutic implications. *J Dermatol* 41:205–12
- Yamaguchi J, Nakamura F, Aihara M et al. (2008) Semaphorin3A alleviates skin lesions and scratching behavior in NC/Nga mice, an atopic dermatitis model. *J Invest Dermatol* 128:2842–9
- Yosipovitch G, Goon A, Wee J et al. (2000) The prevalence and clinical characteristics of pruritus among patients with extensive psoriasis. *Br J Dermatol* 143:969–73

IL-10 Signaling in Dendritic Cells Attenuates Anti-*Leishmania major* Immunity without Affecting Protective Memory Responses

Journal of Investigative Dermatology (2015) **135**, 2890–2894; doi:10.1038/jid.2015.236; published online 16 July 2015

TO THE EDITOR

Leishmania major (*L. major*) parasites cause skin lesions and are transmitted by sand flies infecting millions of people

each year. Upon infection, the parasites (promastigotes) are phagocytosed by macrophages where they transform into amastigotes and replicate. The

amastigotes are eventually released and taken up by dendritic cells (DCs), which activate naive T cells in skin-draining lymph nodes (sdLNs). In *L. major*-infected humans and mice on a resistant background, DCs induce an IFN γ -driven T-helper (Th) type-1/T cytotoxic (Tc) 1 response that stimulates

Abbreviations: DC, dendritic cell; IFN, interferon; IL, interleukin; *L. major*, *Leishmania major*; sdLN, skin-draining lymph node; Th, T helper; Treg, regulatory T cell

Accepted article preview online 22 June 2015; published online 16 July 2015